

ΕΥΠΕΤΗΡΙΟ ΑΓΓΛΙΚΩΝ ΟΡΩΝ

A

Analysis of variance, 293, 743

B

Binomial test, 40

Birnbaum-Hall test, 575

C

Chi-square goodness-of-fit test, 390

Chi-square test,

- for differences in probabilities, 668, 698
- with fixed marginal totals, 688
- for independence, 672

Coefficient

- of concordance, Kendall's, 340

Concordance, Kendall's coefficient of, 340

Confidence band for a distribution function, 448

Confidence interval,

- for the difference between two means, 253, 287
- for the mean, 253
- for the median difference, 253, 287
- for a probability, 119
- for a quantile, 130
- for a slope, 622
- graphical method for, 287

Contingency table, 666, 696

Continuity correction, 709

Correlation,

- sign test for, 108
- rank, 340

Correlation coefficient:

- Kendall's partial, 355
- Kendall's rank, 340, 355
- Pearson's, 361
- Spearman's rank, 326

Correlation, test

- Kendall's rank, 340, 355
- sign test for, 108

Spearman's rank, 326

Cox and Stuart test for trend, 97

Cramér von Mises:

- goodness-of-fit test, 565

D

Difference,

- confidence interval for the median, 253, 287

Difference between two means,

- confidence interval for the, 253, 287

Distribution function,

- confidence band for, 448

E

Exact test, Fisher's, 698

Exponential distribution,

- Lilliefors test for the, 491

F

Families of distributions,

- goodness-of-fit tests for, 470

Fisher's exact test, 698

G

Goodness-of-fit test:

- chi-square, 390
- Cramér-von Mises, 565
- Kolmogorov, 422

Goodness-of-fit tests for families of distributions, 470

H

Hotelling-Pabst test, 337

I

Independence, the chi-square test for, 672

K

k -sample Smirnov test, one sided, 582

- two sided, 592
- K**
 - Kendall's coefficient of concordance, 340
 - Kendall's partial correlation coefficient, 355
 - Kendall's rank correlation test, 355
 - Kendall's tau, 340
 - Kolmogorov goodness-of-fit test, 422
 - Kolmogorov-Smirnov tests, 548
 - Kruskal-Wallis test, 292
- L**
 - Least squares method, 609
 - Lilliefors test for normality, 472
 - for the exponential distribution, 491
 - Limits, tolerance, 169
 - Linear regression, 606, 608
- M**
 - Mann-Whitney test, 261
 - Marginal totals,
 - chi-square test with fixed, 688
 - McNemar test, 722
 - Measures of dependence, 323
 - Median difference,
 - confidence interval for, 253, 287
 - Median test, 733
 - Monotonic regression, 638
 - Moore-Wallis test, 86, 515
- N**
 - Normality:
 - Lilliefors test for, 472
 - Shapiro-Wilk test for, 499
 - One-sided k -sample Smirnov test, 582
- P**
 - Partial correlation coefficient, Kendall's 355
 - Pearson's correlation coefficient, 361
- Q**
 - Quantile,
 - confidence interval for, 119
- R**
 - Randomness, 513
 - Moore-Wallis test for, 515
 - Runs test for, 520
 - Rank correlation, 340
 - Kendall's test for, 340, 355
 - Spearman's test for, 326
 - Regression, 606
 - least squares method, 609
 - linear, 608
 - monotonic, 638
 - slope,
 - confidence interval for, 622
 - test for, 609, 622
 - Theil's method, 622
 - a variant of Theil's method, 623
 - Rho, Spearman's, 326
 - Runs test, 520
- S**
 - Shapiro-Wilk test for normality, 499
 - Siegel-Tukey test, 304
 - Signed ranks test, Wilcoxon, 209, 236
 - Sign test, 70, 86
 - for correlation, 108
 - Smirnov test, 548
 - one-sided k -sample, 582
 - two-sided k -sample, 592
 - Spearman's rank correlation test, 326
 - Spearman's rho, 326
 - Squared ranks test for variances, 311
 - Symmetric distributions, 205, 206
- T**
 - Table, contingency, 666
 - Tau, Kendall's, 340
 - Theil's method, 622
 - a variant of, 623
 - Tolerance limits, 169
 - Trend, Cox and Stuart test for, 97
 - Two-sample Cramér-von Mises test, 565
 - Two-sided k -sample Smirnov test, 592
 - Two-way contingency table, 666

V

Variance,

- analysis of, 293, 743
- Siegel-Tukey test for, 304
- squared ranks test for, 311
- tests for, 304

W

Wilcoxon, test

- signed ranks test, 209, 236
- two-sample test, 203, 261

Wilcoxon Mann-Whitney test, 261